HYDRO-GEOCHEMICAL CHARACTERISTICS OF SURFACE AND SUB-SURFACE WATER OF LALMATIA COAL MINE AREA, RAJMAHAL BASIN, DISTRICT- GODDA (JHARKHAND), INDIA

A Synopsis submitted to
University Department of Geology,
Ranchi University, Ranchi

For the degree of
DOCTOR OF PHILOSOPHY (Ph.D.)

Submitted by:
SULEKHA MURMU
Research Scholar

Under the supervision of :
Dr. CHANCHAL LAKRA
Assistant Professor
University Department of Geology
Ranchi University

The topic of the research :

HYDRO-GEOCHEMICAL CHARACTERISTICS OF SURFACE AND SUB-SURFACE WATER OF LALMATIA COAL MINE AREA, RAJMAHAL BASIN, DISTRICT-GODDA (JHARKHAND), INDIA

1. A BRIEF INTRODUCTION OF THE RESEARCH TOPIC

Water is an extremely valuable natural resource that is used extensively for home, commercial, industrial, and agricultural purposes worldwide and that enters the environment through a variety of sources [1,2,3].

The quantity and quality of water are equally significant [4,5]. But in places where there is a mining and mineral processing industry, the problem of poor water quality is especially severe. Coal mining plays a significant role in global economic development of many nations [6,7,8]. India currently ranks third in the world for coal production, with over 319 billion tonnes of coal reserves [9,10]. The most identifiable and observable environmental issue is caused by coal mining because the mining process changes the physical, chemical, and biological parameters of the surrounding environment and has a significant impact on both the ecological unit and human civilization [1,11,3]. Particularly during mining operations, a number of waste classes are created, which have the potential to produce different kinds of pollution before contamination happens [1,12,3].

Mining can negatively impact the quantity and quality of groundwater and surface water in mining areas [13,14] endangering the water resources' sustainability [15] and general health of the local population [16,17]. There have been reports of ecological deterioration and health issues caused by mine waste spills all over the world [18,19,20,21]. So, it is of great importance to evaluate the ground water in mining areas hence attracting attention from both policy makers and researchers [21].

The knowledge on hydro-chemical and geochemical characteristics is more crucial to evaluate the quality of surfacewater and groundwater in order to determine its suitability for various uses [3].

Groundwater quality along with hydrogeochemical studies in coal mining areas is essential for comprehending the working mechanisms and processes because it includes resource exploitation, overburden removal, the production of coal mine waste and mine drainage, and the offsetting of the equilibrium in the local and regional aquatic systems. Hydrochemical characteristics give essential details regarding the sources, migration, circulation, and recharge of groundwater [22,10].

2. STATE OF THE ART

Natural processes like rock-water interaction and seawater intrusion have an impact on groundwater quality[23], in addition to manmade inputs like those from mining activities [24]. The hydrogeological context of active coal mining areas, combined with large overburden (OB) material dumps, presents a challenging set of obstacles for quantifying and comprehending the mechanisms that regulate water quality. These human-caused interventions can occasionally start processes and interactions that seriously harm groundwater resources [25]. When coal is extracted from open-pit mines, waste or OB materials are created. Some of these materials are discarded inside the mine as internal dumps, and others are piled onto the surface next to the

mines as external dumps. Water that has precipitated penetrates through these OB materials and into the nearby groundwater aquifer through active strip mines. After being used for mining operations, excess water from mines is either released onto the surface for irrigation or used for residential use [26]. Therefore, unless safety precautions are taken, there could be dangerous outcomes. Furthermore, the water table is significantly lowered as a result of mine dewatering, which is necessary for all opencast and underground activities [27,28]. This drop in the water table also causes groundwater from nearby places that may include pollutants and dissolved materials to flow into the earth, which has an impact on the groundwater's quality. As a result, having accurate information about water quality is essential since it affects how well water resources are managed and used [29,30,31].

2.1 LOCATION OF THE STUDY AREA

Lalmatia coal mine is located in godda district which comes under the Boarijor CD block in godda subdivision in the Indian state of Jharkhand. The block is located between latitudes $25^0 1'12''$ N and $25^0 3'15''$ N and Longitude $87^0 21'0''$ E and $87^0 24'0''$ E and occupies an area of around 15 sq. km. The surrounding villages are Lalmatia, Dakaita, Hijukita, Basdiha, Lohandia, Bhorai, Bara Simra, Neema, Kendua, Telgama, Kusuma, Tetaria, Ballia etc.



Fig(a):map of India(insmandovi.kvs).



Fig(b):map of Jharkhand (tourism.jharkhand.gov.in).



Fig(c): Geological map of Lalmatia coal mine (35).

Fig 1: Location map of the study area.

The study area falls in the Survey of India Toposheet no. G45P8 (OSM no.) and 72O/8 (sheet no.) (scale 1:50,000).



Fig 2 : Toposheet map of the study area (Lalmatia, district- Godda),Jharkhand.

OSM no.-G45P8, Sheet no.- 72O/8 (Scale 1:50,000) (Survey of India).

2.2 GEOLOGY AND STRATIGRAPHIC SEQUENCE

The Rajmahal Basin is thought to have developed from tholeiitic magmatism during the Gondwana period. The Barakar and Talchir formations are situated beneath the undifferentiated Triassic layers beneath the trap [32]. In the foot of the Rajmahal hills, the lower Gondwana exposures comprise the Rajmahal coal fields, which are generally aligned in a north-south direction. There is a depositional contact between the Barakar Formation and the Archeans that is located toward the west. Parts of the coalfield have also been found to expose the lithostratigraphic units Dubrajpur, which underlies the Rajmahal traps, and Talchir, which underlies the Barakar formation.The general geological succession exposed in the Rajmahal area, eastern Jharkhand is given in Table 1 [33,34].

The Lalmatia exploration block, including 15 square kilometers, is home to the Rajmahal open cast mine and has been explored by the Central Mine Planning and Design Institute (CMPDI) [35].

Table 1: Geological succession exposed at the Rajmahal area, eastern Jharkhand (after Raja Rao, 1987[33]; Singh and Singh, 1996[34]).

Age	Formation	Lithology	Thickness
Recent to sub-recent	Alluvium	Loose soil, silt and clay	Upto 80 m
----------UNCONFORMITY----------			
Early Cretaceous to Early Jurassic	Raajmahal Traps and Inter-trappeans	Flows of basalt, pitchstone and Inter-trappean beds (sandstone, shale and ash)	600 m
----------UNCONFORMITY----------			
Late Triassic	Dubrajpur	Pebbly sandstone, Coarse to medium grained sandstone, red siltstone	60-250 m
----------UNCONFORMITY----------			
Early Permian	Barakar	Coarse to medium grained sandstone, pebbly sandstone, gray shales, clay and coal seams	250-550 m
	Karharbari	Grits, conglomerates and carbonaceous sandstones	
	Talchir	Tillite, fine to medium grained sandstones, olive green shale	
----------UNCONFORMITY----------			
Archean		Basic rocks, amphibolites, bands of quartzite, limestone, gneisses and granite	

2.3 HYDROGEOLOGY

Godda district enjoys a cold winter, a good monsoon, and an oppressively hot and dry summer. The year can be divided into four seasons. Summer extends from March through the first week of June, and winter spans from December to February. The months of the southwest monsoon are June through September. The post-monsoon season, which spans from October to November, occurs between the monsoon and winter seasons. The highest rainfall occurs in July [36].

In all of the lithological units that make up the shallow aquifers, ground water is primarily found in a phreatic state. In deeper aquifers, it is occasionally found in semiconfined and confined conditions [37].



LEGEND

AGE GROUP	LITHOLOGY	SYMBOL	HYDROGEOLOGICAL CONDITION	GROUND WATER POTENTIAL
			UNCONSOLIDATED / POROUS FORMATION	
Quaternary	Recent & older alluvium clay, sand, gravel, calcareous & ferruginous concretion		Moderatelly thick, regionally extensive unconfined / semi confined aquifer down to 100m.	Good yield prospect 25 - 75 m ³ / hr.
Tertiary	Laterite		Thin veneer in patches under unconfined aquifer down to 10 - 15 m.	Limited yield prospect 5 m ³ / hr.
			SEMICONSOLIDATED FORMATION	
Gondwana super group	Sandstone, shale, siltstone, with coal seams.		Moderatelly thick, regionally extensive confined qquifer / free flowing condition down to 200m.	Very good yield prospect 50 - 100 m³ / hr.
			CONSOLIDATED FORMATION / FISSURED FORMATION	
Rajmahal Trap	Basalt with intertrappean sandstones & shales		Groundwater restricted to weathered zone, fractured zone. Contact of two flows & intertrappeans down to 100m.	Limited yield prospect blow 25 m ³ / hr at selective places.
Precambbrian	Granite - gneiss, mica schist amphibolotes etc.		Groundwater restricted to weathered zone, fractured zone. down to 150m.	Moderately good yield prospect 10 - 40 m ³ / hr at selective places.

Fig 3 : Hydrogeology of the godda district [37].

2.4 DRAINAGE

The district's main rivers are the Harna, Sunder, Sapin, Kao, Cheer, and Geura. The drainage generally flows in a SE-NW direction. The structural elements, in particular the joints and foliation, have a significant influence on drainage and regulate the district's drainage pattern [37].

2.5 LITERATURE REVIEW

Ball (1877) studied the structure and stratigraphy of Rajmahal Basin [38]. Exploration was done in this area by numerous workers (Madabhushi and Mukherjee, 1980[39]; Roy et al., 1983[40]; Das et al., 1985[41]; Madabhushi, 1985[42a], 1990[42b]; Singh, 1992[43]). The Coal seams are underlain by the Barakar Formation is reported by Raja Rao (1987)[33]. Few reports exist on the organic petrography and geochemical characterization of coal by Singh (1992)[43] and Singh and Singh (1996)[34]. Due to the abundance of plant megafossils and palynomorphs, the intertrappean beds of Rajmahal have been thoroughly investigated (Sahni, 1932[44a], 1948[44b]; Sahni and Rao, 1933[45]; Rao, 1943[46]; Vishnu-Mittre,1956[47a], 1958[47b]; Sharma, 1967[48a], 1969[48b], 1974[48c], 1997[48d]; Sengupta, 1988[49]; Banerjee, 1993[50a], 1995[50b], 2000[50c]; Banerji and Jana, 1998[51]; Tiwari et al., 1984[52]; Baksi et al., 1992[53]; Tiwari and Tripathi, 1995[54]; Tripathi, 2008[55])[35]. On the coal-bearing sequences in the Rajmahal Basin, numerous investigations have been conducted particularly on the morpho-taxonomical (Feistmantel, 1880[56]; Singh et al., 1987[39]; Bajpai and Maheshwari, 1991[57]; Maheshwari and Bajpai, 1992[58]; Srivastava and Pant, 2002[69]), palynology (Tripathi and Ray, 2005[60]; Tripathi et al., 2010[61]), petrological (Roy et al., 1983[40]; Madabhushi, 1990[42b]; Singh, 1992[43]), molecular composition and hydro-carbon potential (Tiwari et al., 2016[62a], 2017[62b]) aspects [32].

Work on ground water resources have been assessed by Remote sensing technology (Saraf & Choudhary (1998)[63]; Asadi et al. (2007)[64]; Singh (2007)[65]; Latha (2010)[66]; Singh and Khan (2011)[67]; Ilayaraja and Ambika (2015)[68]; Oseke et al. (2021)[69]. The work on A correlational study of uranium in groundwater with other physicochemical parameters using GIS mapping

in Godda district of Jharkhand, India was done by Srivastava et. al., (2023) [36]. Other work is on Geochemical mobilization of Arsenic, Chromium and Uranium in Gangetic Plain of Bihar and Jharkhand done by Jha et. al., (2023) [70].

3. PROBLEMS TO BE INVESTIGATED

- To estimate the quality of ground water and surface water.
- To estimate the water quality index (WQI).
- To examine the groundwater parameters.
- Characterization of hydro-geochemical facies variations.
- Assessment of various factors that control the quality of groundwater, river water and their suitability for various utilization.
- Investigation of the effect of active coal mining on groundwater resources of the study area.
- Its impact on public health and the extent to which water can be used in various ways based on its quality.

4. OBJECTIVES OF THE INVESTIGATION AND MOTIVATION

- To understand the water quality and ecology of the area affected by coal mining.
- To understand the effect of mining on the natural resources utilised.
- To make a database of groundwater chemistry in the coal mining areas, which will be useful in future assessment for the impact of coal mining operations in the study area.

Motivation : In coal mining locations, common issues include the deterioration of water quality, the production of acidic mine drainage water, the discharge of heavy metals, and mixing with nearby aquifers and water supplies. All of these factors have a major negative influence on groundwater resources. In order to maintain drinking water standards for both the present and the future, it is imperative that water quality indicators be regularly monitored and assessed [10].

5. RESEARCH HYPOTHESIS

The hydrogeochemical characteristics of the groundwater provide a critical framework for assessing the quality of groundwater and the aquifer's solute acquisition process.The number of hydrogeochemical processes that result in a change in the fluid medium's chemical composition and do not conflict with one another control the flow of groundwater throughout its course from recharge to discharge locations. Therefore, understanding the hydrogeological system and the processes involved in the hydrogeochemical evolution is essential to designing a sustainable ecosystem[71,72]. In order to evaluate surface and groundwater sources of drinking water, a thorough impact assessment of the principal ions in the physicochemical parameters must be conducted. As a result, an indexing technique called the water quality index (WQI) offers a cumulative evaluation of drinking water quality criteria [10].

The investigation on the hydrogeochemical characteristics of ground water and surface water of the Lalmatia coal mine area has not been done yet by anyone. So, it is a virgin area and for future investigation I have opted for the topic:- hydro-geochemical characteristics of surface and sub-surface water of Lalmatia coal mine area.

6. RESEARCH DESIGN AND METHODOLOGY

The research design and methodology will encompass:

Step 1:

- Literature survey
- Field survey
- Sample collection pre monsoon and post monsoon
- Photography
- Classification
- Make arrangement for storage of the sample

Step 2 : Research analysis will be carried out in the following stages:

- Chemical analysis
- Evaluation of water quality index (WQI)
- Health risk assessment
- Interpretation and correlation

❖ **Collection of samples:**

- The samples of groundwater will be collected from dug wells, bore wells and hand pump and the surface water samples from ponds, rivers and mine effluent water in the study area.
- All samples will be collected in a 1-litre capacity polyethylene bottles. Before sampling, the bottles will be washed with respective samples to avoid being contaminated.
- After tapping the water, in situ parameters such as hydrogen ion concentration (pH), temperature (T), total dissolved solids (TDS), and electrical conductivity (EC) will be determined at the sampling sites itself with the help of a pH meter and a portable multi-parameter tool kit.

❖ **Chemical analysis**: The physicochemical analysis of water samples will be carried out as per the procedure using standard procedure suggested either by Indian standard (IS) or American Public Health Association (APHA 2017).

❖ **Evaluation of Water Quality Index (WQI) :**

- WQI of water sample will be calculated by combining the values of several parameters to produce a single figure that represents the water's quality or level of pollution [73].
- Different weights will be assigned for physicochemical parameters .

❖ **Interpretation and correlation** : The interpretation and correlation will be made on the basis of results obtained by above analysis.

7. SOURCES OF DATA AND TOOLS TO BE USED IN COLLECTION OF DATA

- Clinometer compass
- Handheld GPS
- Toposheet
- Markers
- 1 litre capacity Polyethylene bottles
- Notepad and Pen
- Camera
- pH meter
- a portable multi-parameter tool kit

8. SIGNIFICANCE OF THE STUDY

- The groundwater chemistry study will be useful to study the impact of coal mining on the water resources of the area.
- The calculation of the water quality index (WQI) determines whether the quality of water is fit for drinking purposes.
- Health risk assessment is a crucial prerequisite for studies relating to groundwater quality appraisal. It is a set of techniques that are aimed at evaluating the potentially harmful impact of groundwater on human health [74,75,76,77,78,79,10].

9. ITS RALEVANCE TO THE PRESENT DAY PROBLEM AND NEED TO THE SOCIETY

Water quality and chemistry are impacted by coal mining operations both during the active mine life and even decades after the mine closes [80,81,10]. One of the main things that can lower the rates of baby and mother mortality is clean drinking water. Long-term exposure to pollutants is harmful to people's health and well-being. High levels of pollution in water are extremely dangerous for human health and endanger the entire environment that supports life.

10. CONTRIBUTION TO THE EXISTING KNOWLEDGE

Policymakers and water management boards will benefit from the research's findings in terms of access to clean drinking water. The understanding of the hydrological milieu and the mechanisms influencing the groundwater chemistry in the coal mining sites may be greatly enhanced by this investigation. For the purpose of monitoring, long-term protection, effective management of water resources, and preventing health concerns associated with coal mining operations in the area, a baseline database on water chemistry will be generated.

11. FUTURE WORK

- The current study's findings will support the efforts of health authorities to conduct additional epidemiological investigations and gather comprehensive data regarding human exposure to ions found in surface and ground water.
- Managing the precious water resource and meeting future water demands in the coalfield area will be made much easier with the baseline data on water analysis.
- The resulting data will be helpful to the NGO's (national or international) working for water quality and its impact on living beings.
- On the basis of obtained data, a detailed elemental research work can be initiated for that particular area and also for other mining areas.
- This paper will be helpful to the researchers for their study in that area in future.
- This study of the water analysis can be helpful for the students and research scholars of geology, geography and environmental science.

12. RELATED REFERENCES

1) Howladar, M. F., Deb, P. K., Muzemder, A. T. M. S. H. & Ahmed, M. (2014) : Evaluation of water resources around Barapukuria coal mine industrial area, Dinajpur, Bangladesh. Appl Water Sci 4: DOI 10.1007/s13201-014-0207-5., pp. 203-222.
2) Seddique, A. A. and Matin, K. A. (2013) : Assessment of groundwater vulnerability in and around Narayanganj town, Bangladesh : Insight into Groundwater contamination. Intern Jour of Env, Vol.3 (1), pp. 10-18.
3) Khan, Md. H. R., Seddique, A. A., Rahman, A., Shimizu, Y. (2017) : Heavy Metals Contamination Assessment of Water and Soils in and Around Barapukuria Coal Mine Area, Bangladesh. Amer. Journ. of Env. Protec. Vol. 6, No. 4, pp.80-86.
4) Todd, O. K. (1959) : Ground Water Hydrogeology. Reprint Edition, John Wiley and Sons Inc. New York. Chandra, C. 1992 Jharia Coalfield, Geo. Soci. of India, Bangalore, pp. 1-11.
5) Gupta, S. K., Nikhil, K. (2016) : Ground Water Contamination in Coal Mining Areas: A Critical Review. Inter. J. of Engi and Appl. Sci. (IJEAS). ISSN: 2394-3661, Volume-3, Issue-2.
6) Schneider, L., Maher, W. A., Rose, NL et al (2020) : Assessing environmental contamination from metal emission and relevant regulations in majior areas of coal mining and electricity generation in Australia. Sci Total Environ 728:137398.
7) Li, P., Qian, H., Howard, K. W. F., Wu, J. (2015) : Building a new and sustainable " Silk Road economic belt". Environ Earth Sci., pp.7267-7270.
8) Wang, S., Tang, H., Shi, L., Liu, J., Yang, Z., Zhu, H., Xu, F., Zhu, K., Fan, J., Fang, G. (2022) : Hydrogeochemical analysis and assessment of mine water quality in Tangjiahui mining area, Inner Mongolia, China : Environ Earth Sci 84:49.
9) GSI. (2019). Indian Coal and Lignite Resources-2019. GSI, Natur. Ener. Resour. Mission-II B, Gov. Of India, pp. 47.
10) Samal, P., Mohanty, A. K., Khaoash, S., Mishra, P. (2022) : Hydrogeochemical Evaluation, Groundwater Quality Appraisal, and Potential Health Risk Assessment in a coal Mining Region of Eastern India : Water Air Soil Pollut 233:324.

11) Halim, M. A., Majumder, R. K., Zaman, M. N., Hossain, S., Rasul, M. G. & Sasaki, K. (2012) : Mobility and impact of trace metals in Barapukuria coal mining area, Northwest Bangladesh., Springer., Arab J Geosci DOI 10.1007/s12517-012-0769-1., pp.13.
12) Farhaduzzaman, M., Abdullah, W. H. And Islam, M.A. (2012) : Depositional environment and hydrocarbon source potential of the Permian Gondwana coals from the Barapukuria basin, Northwest Bangladesh. Intern J. of Coal Geo., Vol. 90-91, No. 1, pp. 162-179.
13) Bebbington, A., Williams, M. (2008) : Water and mining conflicts in Peru. Mt. Res. Dev 28(3-4):190-195.
14) Vidic, R. D., Brantley, S. L., Vendenbossche, J. M., Yoxtheimer, D., Abad, J. D. (2013) : Impact of shale gas development on regional water quality, Science 340(6134):1235009
15) Yang, H., Flower, R. J., Thompson, J. R. (2013) : Shale-gas plans threaten China's water resources. Science 14340(6138):1288.
16) Li, P., Li, X., Meng, X., Li, M., Zhang, Y. (2016) : Appraising groundwater quality and health risks from contamination in a semiarid region of northwest China. Expo Health 8(3): 361-379
17) Wu, J., Sun, Z. (2016) : Evaluation of shallow groundwater contamination and associated human health risk in an alluvial plain impacted by agricultural and industrial activities, mid-west China. Expo Health 8(3):311-329
18) Esteller, M. V., Dominguez-Mariani, E., Garrido, S. E., Aviles, M. (2015) : Groundwater pollution by arsenic and other toxic elements in an abandoned silver mine, Mexico. Environ Earth Sci 74(4):2893-2906
19) Njinga, R. L., Tshivhase, V.M. (2017) : Major chemical carcinogens in drinking water sources : health implications due to illegal gold mining activities in Zamfara State-Nigeria. Expo Health. https://doi.org/10.1007/s12403-017-0265-7
20) Wright, I. A., Belmer, N., Davies. P. J. (2017) : Coal mine water pollution and ecological impairment of one of Australia's most 'protected' high conservation-value rivers. Water Air Soil Poll 228:90. https://doi.org/10.1007/s11270-017-3278-8
21) Li, P. (2018) : Mine water Problems and Solutions in China. Mine water and Envi. 37. Pp. 217-221
22) Selfie, A., Aris, A.Z., Ramli, M. F., Narany, T. S., Shamsuddin, M. K. N., Saadudin, S. B. & Zali, M. A. (2018) : Hydro-geochemistry and groundwater quality assessment of the multilayered aquifer in Lower Kelantan Basin. Malaysia Env. Earth Sci., 77(10), 397.
23) Sophiya, M. S., Syed, T. H. (2013) : Assessment of vulnerability to seawater intrusion and potential remedial measures for coastal aquifers : a case study from eastern India. Environ. Earth Sci., pp.1197-1209.
24) Kurdi, M., Hezarkhani, A., Eslamkish, T. (2014) : Assessment of chemical properties and hydro-geochemical coefficients at the QarehSou Basin, Golestan Province, Iran. Environ. Earth Sci., v.72, pp.3243-3249
25) Bukowski, P., Bromek, T., Niak, A. I. (2006) : Using the DRASTIC system to assess the vulnerability of ground water to pollution in mined areas of the upper Silesian Coal Basin. Mine Water Environ., v.25, pp.15-22

26) Soni, A. K., Wolkersdorfer, C. (2015) : Mine water : policy perspective for improving water management in the mining environment with respect to developing economies. Int. J. Min. Reclam. Environ., Doi:10.1080/17480930.2015.1011372
27) Adhikari, K., Sadhu, K., Chakroborty, B., Gangopadhyay, A. (2013) : Effect of mining on geochemistry of groundwater in Permocarboniferous Gondwana coalfields: Raniganj Basin, India. J. Geol. Soc. India, V.82, pp.392-402
28) Singh, R. B., Doley, T., Chakraborty, T., Mohanty, A. K., Kumar, M. (2015) : Coal mining impact assessment on groundwater system by virtue of piezometer installation: a case study from Kusmunda OC project of SECL, Chhattisgarh, Central India. Min. Tech., v.36(1), pp.29-35
29) Reddy, A. G. S. (2013) : Evaluation of hydrogeochemical characteristics of phreatic alluvial aquifers in southeastern coastal belt of Prakasam district, South India. Environ. Earth Sci., v.68, pp.471-485.
30) Pereira, A. J. S. C., Pereira, M. D., Neves, L. J. P. F., Azevedo, J. M. M., Campos, A. B. A. (2015) : Evaluation of groundwater quality based on radiological and hydrochemical data from two uriniferous regions of western Iberia: Nisa (Portugal) and Ciudad Rodrigo (Spain). Environ. Earth Sci., v.73, pp.2717-2731.
31) Singh, R., Syed, T. H., Kumar, S., Kumar, M., Venkatesh, A. S. (2017) : Hydrogeochemical assessment of surface and groundwater resources of Korba coalfield, Central India: environmental implications. Arab. J. Geosci., v.10, pp.318.
32) Mathews, R. P., Pillai, S. S. K., Manoj, M. C., Agrawal, S. (2020) : Palaeoenvironmental reconstruction and evidence of marine influence in Permian coal-bearing sequence from Lalmatia coal mine (Rajmahal Basin), Jharkhand, India: A multi-proxy approach : Intern. J. of Coal Geol., v.224, pp.103485.
33) Raja Rao, C. S. (1987) : Coal resources of Bihar (excluding Dhanbad district). Bull. Geol. Surv. India Series, V. 45, pp.300-322.
34) Singh , K. J. and Chandra, S. (1996) : Plant fossils from the exposure near Gopal Prasad Village, Talcher Coalfield, Orissa, with remarks on the age of the bed. Geophytology, v.26, pp.69-75.
35) Pillai, S. S. K., Mathews, R. P., Murthy, S., Goswami, S., Agrawal, S., Sahoo, M. and Singh, R. K. (2020) : Palaeofloral Investigation Based on Morphotaxonomy, Palynomorphs, Biomakers and Stable Isotope from Lalmatia Coal Mine of Rajmahal Lower Gondwana Basin, Godda District, Jharkhand : An Inclusive Empirical Study : Jour. Geol. Soc. of India.
36) Srivastava, M. and Srivastava, P. K. (2023) : A correlational study of uranium in groundwater with other physicochemical parameters using GIS mapping in Godda district of Jharkhand, India : Environ. Geochem. Health, v.45, pp.9903-9924.
37) Central Ground Water Board (2013) : Ground Water Information Booklet, Godda district, Jharkhand State.
38) Ball, V. (1877) : On the geology of the Mahanadi Basin and its vicinity. Rec. Geol. Surv. India, v.10, pp.167-185.
39) Madabhusi, S. and Mukherjee, G. (1980) : A report on the geological mapping in the North Extension of Rajmahal Coalfield, Bhagalpur and Santhal Parganas District, Bihar. Unpubld. G.S.I. Prog. Report.

40) Roy, S., Mukherjee, A. K. and Madabhushi, S. (1983) : Report on regional exploration for coal in Hura block, Rajmahal Coalfield, Santhal Parganas, Bihar. Geol. Surv. India. Unpbld. Prog. Rept.
41) Das, A. K., Bandopadhyay, D. N. and Madabhushi, S. (1985) : Regional exploration for coal in Northern extension area of Rajmahal Group of coalfields. Geol. Surv. India, Unpubl. Prog. Rept., pp.1980-1984.
42) (a) Madabhushi, S. (1985) : Structure, lithofacies, sedimentary tectonics and coal potentials of Rajmahal Hills, Bihar, India. Unpub. Ph.D. thesis.
(b) Madabhushi, S. (1990) : Linear ash moisture relation of coal seam in a Borehole drilled in Chuoerbhita Coalfield, Bihar. J. Geol. Soc. India, v.35, pp.644-648.
43) Singh, P. K. (1992) : Organic petrology and geochemical characterisation of the Permian coal deposit of the Rajmahal Basin, Bihar. Unpub. Ph.D. thesis.
44) (a) Sahni, B. (1932) : A petrified Williamsonia (w. sewardiana sp. nov.) from the Rajmahal Hills. Mem. Geol. Surv. India. Palaeonl. indica,n.set. 20, pp.1-19.
(b) Sahni, B. (1948) : The Pentoxyleae: a new group of Jurassic gymnosperms from the Rajmahal Hills of India. Bot. Gaz., v.110. pp.47-80.
45) Sahni, B. and Rao, A. R. (1933) : On some Jurassic plants from the Rajmahal Hills. Jour. Asiatic Soc. Bengal, New Series, v.27, pp.183-208.
46) Rao, A. R. (1943) : Jurassic spores and sporangia from the Rajmahal Hills, Bihar. Proc. Nat. Scad. Sci., India (B), v.13, pp.181-197.
47) (a) Vishnu-Mittre (1956) : Masculostrobus sahni sp. nov., a petrified conifer male cone producing three-winged and one-and four-winged abnormal pollen grains from the Jurassic of the Rajmahal Hills, Bihar. Grana. Palynol., v.1, pp.99-107.
(b) Vishnu-Mittre (1958) : Studies on the fossil flora of the Nipania, Rajmahal series, India-Pteridophyte and general observations on Nipania fossil flora. Palaeobotanist, v.7, pp.47-66.
48) (a) Sharma, B. D. (1967) : Investigations on the Jurassic flora of Rajmahal Hills, India, 3. A review of the genus Ptilophyllum of Morris, with description of two new species from Amarjola in the Rajmahal Hills. Palaeontogr. Abt. B, v.120, pp.139-150
(b) Sharma, B. D. (1969) : On a collection of fossil fern from Dhokuti in the Rajmahal Hills, India. Palaeontogr. Abt. B, v.128, pp.55-63.
(c) Sharma, B. D. (1974) : Ovule ontogeny in Williamsonia Carr. Palaeotogr. Abt. B, v.148, pp.137-143.
(d) Sharma, B. D. (1997) : An early angiosperm fructification resembling Lesqueria Crane and Dilcher from the Rajmahal Hills, India. Phytomorphology, v.47, pp.305-310.
49) Sengupta, S. (1988) : Upper Gondwana stratigraphy and palaeobotany of Rajmahal Hills, Bihar, India. Mem. Geol. Surv. India. Palaeont. Indica, v.48, pp.1-180.
50) (a) Banerji, J. (1993) : Plant fossils from Chunakhal, Rajmahal Hilss, Bihar. Geophytology, v.23, pp.21-80.
(b) Banerji, J. (1995) : Recent records of Mesozoic ferns from Rajmahal basin. Indian Fern J., v.12, pp.70-78.
(c) Banerji, J. (2000) : Megafloral diversity of the upper Gondwana sequence of the Rajmahal Basin, India. Jour. Africa Earth Sci., v.31, pp.133-144.
51) Banerji, J. and Jana, B. N. (1998) : Early Cretaceous megafossils from Balidih, Rajmahal basin, India. Geophytology, v.27, pp.33-36.

52) Tewari, R. S., Kumar, P. and Tripathi, A. (1984) : Palynodating of Dubrajpur and Intertrappean beds in subsurface strata of northeastern Rajmahal Basin.In: Proceeding V Indian Geophytological Conference, Lucknow 1983, Special Publication, pp.207-225.
53) Baksi, A. K., Tiwari, R. S., Tripathi, A. and Farrar, E. (1992) : Geochemical, geochronological and palynological observations on Early Cretaceous lavas in the Rajmahal Basin, p. 16-17.
54) Tewari, R. S. and Tripathi, A. (1995) : Palynological assemblages and absolute age relationship of intertrappean beds in the Rajmahal Basin, India. Cretaceous Res., v.16, pp.53-72.
55) Tripathi, A. (2008) : Palynochronology of Lower Cretaceous volcano-sedimentary succession of the Rajmahal Formation in the Rajmahal Basin, India. Cretaceous Res., v.29, pp.913-924.
56) Feistmantel, O. (1880) : The fossil flora of Gondwana system. The flora of Damuda-Panchet Divisions. Mem. Geol. Surv. India. Palaeont. Indica, Series, v.12, pp.1-149.
57) Bajpai, U. and Maheshwari, H. K. (1991) : On two enigmatic infructescences from Permian Gondwana of the Rajmahal Basin. Palaeobotanist, v.39, pp.9-19.
58) Maheshwari, H. K. and Bajpai, U. (1992) : Ginkgophyte leaves from Permian Gondwana of Rajmahal Basin, India. Palaeontogr. Abt. B, v.224, pp.131-149.
59) Srivastava, P. C. and Pant, D. D. (2002) : Some new pteridophytic remains from the Permian strata of Rajmahal Hills, India. Palaeobotanists, v.51, pp.45-55.
60) Tripathi, A., Ray, A. (2005) : Palynodating of coal bearing sediments in Pachwara Coalfield, Rajmahal Basin, Santhal Parganas, Bihar. GSI spcl. Publ., v.81, pp.135-142.
61) Tripathi, A., Murthy, S., Singh, R. K. (2010) : Palynodating of coal-bearing strata near kunda Pahari, Pachwara Coalfield, Rajmahal Basin, Jharkhand, India. J. Palaeontol. Soc. Ind., v.55, pp.29-35.
62) (a) Tewari, A., Dutta, S., Sarkar, T. (2016) : Organic geochemical characterization and shale gas potential of Permian Barren Measures Formation, West Bokaro Basin, India. J. Pet. Geol., v.39, pp.49-66.
(b) Tewari, A., Dutta, S., Sarkar, T. (2017) : Biomarker signatures of Permian Gondwana coals from India and their palaeobotanical significance. Palaeogeogr. Palaeoclimtl. Palaeocol., v.468, pp.414-426.
63) Saraf, A. K., & Choudhary, P. R. (1998) : Integrated remote sensing and GIS for groundwater exploration and identification of artificial recharge sites. Intern. J. of Remt. Sens., v.10, pp.1825-1841.
64) Asadi, S. S., Vuppala, P., & Reddy, A. M. (2007) : Remote sensing and GIS technique for evaluation of groundwater quality in Municipal Corporation of Hyderabad (Zone-V), India. Intern. J. of Envn. Res. And Pub. Health, v.4(1), pp.45-52.
65) Singh, D. S. H., & Lawrence, J. F. (2007) : Groundwater quality assessment of shallow aquifer using geographical information system in part of Chennai city Tamil Nadu. J. Geol. Soc. Of India, v.69(5), pp.1067-1076.
66) Latha, P. S., & Rao, N. K. (2010) : Assessment and Spatial distribution of quality of groundwater in Zone II and III, Greater Visakhapatnam, India using water quality index(WQI) and GI. Intern. J. of Env. Sci., v.1(2), pp.198-212.
67) Singh, P., & Khan, I. A. (2011) : Groundwater quality assessment of Dhankawadi ward of Pune by using GIS. Intern. J. of Geom. and Geosci., v.2(2), pp.668-703.

68) Ilayaraja, K., & Abica, A. (2015) : Spatial Distribution of groundwater quality between Injambakkam-Thiruvanmyiur areas South-East Coast of India. Nat. Env. And Poll. Tech., an Intrna. Quart. Sci. J., v.14(4), pp.771-776.
69) Oseke, F. I., Anornu, G. K., Adjei, K. A., & Eduvie, M. O. (2021) : Assessment of water quality index in reservoirs affected by water diversion. Wat.-Ener. Nex., v.4, pp.25-34.
70) Jha, A. K., SubhajitSikdar, Verma, S., Sharma, U., Thakur, R., Majumder, S., Anand, A., Kumari, P. (2023) : Geochemical mobilization of Arsenic, Chromium and Uranium in Gangetic Plain of Bihar and Jharkhand. Intrn. J. of Chem. And Bioch. Sci., v.24(6), pp.836-841.
71) Sophocleous, M. (2010) : Understanding and explaining surface tension and capillarity: An introduction to fundamental physics for water professionals. Hydrog. J., pp.811-821.
72) Brkic, Z., Briski, M., & Markovic, T. (2016) : Use of hydrochemistry and isotopes for improving the knowledge of groundwater flow in a semi- confined aquifer system of the Eastern Slavinia (Croatia). Catena, v.142, pp.153-165.
73) Gaur, N., Sarkar, A., Dutta, D., Gogoi, B. J., Dubey, R. & Dwivedi, S. K. (2022) : Evaluation of water quality index and geochemical characteristics of surfacewater from Tawang India. Scientific reports.
74) Wu, J., & Sun, Z. (2016) : Evaluation of shallow groundwater contamination and associated human health risk in an alluvial plain impacted by agricultural and industrial activities, mid-west China. Expo. And Health, v.8(3), pp.311-329.
75) He, S., & Wu, J. (2019) : Hydrogeochemical characteristics, groundwater quality and health risks from hexavalent chromium and nitrate in groundwater of Huanhe formation in Wuqi Country Northwest China. Expo and Health, v.11(2), pp.125-137.
76) Adimalla, N., Li, P., & Qian, H. (2019) : Evaluation of groundwater contamination for fluoride and nitrate in semi-arid region of Nirmal Province, South India: A special emphasis on human health risk assessment. Human and Ecological Risk Assessment: An Intern. J., v.25(5), pp.1107-1124.
77) Prathap, A., & Chakraborty, S. (2019) : Hydrochemical characterization and suitability analysis of groundwater for domestic and irrigation uses in opencast coal mining areas of Chari and Kuju, Jharkhand India. Groundwater for Sustainable Development, 9, 100244.
78) Subba Rao, N., Sunitha, B., Sun, L., Spandana, B. P., & Chaudhary, M. (2020) : Mechanisms controlling groundwater chemistry and assessment of potential health risk: A case study from South India. Geochemistry, 80, 125568.
79) Zhou, M., Li, X., Zhang, M., Liu, B., Zhang, Y., Gao, Y., Ullah, H., Peng, L., He, A., & Yu, H. (2020) : Water quality in a worldwide coal mining city: A scenario in water chemistry and health risks exploration. J. of Geochem. Expl., 213, 1065513.
80) Price, P., & Wright, I. A. (2016) : Water quality impact from the discharge of coal mine wastes to receiving streams: Comparision of impacts from an active mine with a closed mine. Water Air Soil Poll., v.227, pp.155.
81) Tran, T. Q., Banning, A., Wisotzsky, F., & Wohnlich, S. (2020) : Mine water hydrogeochemistry of abandoned coal mines in the outcropped Carboniferous formations, Ruhr area Germany. Environ. Earth Sci., v.79, pp.84.

SIGNATURE OF RESEARCH SCHOLAR

SULEKHA MURMU
Research Scholar
University Dept. Of Geology,
Ranchi University, Ranchi
Jharkhand

SIGNATURE OF SUPERVISOR

Dr. CHANCHAL LAKRA
Assistant Professor
University Dept. Of Geology,
Ranchi University, Ranchi
Jharkhand